


10. Grogan M, Smith HC, Gersh BJ, Wood DL. Left ventricular dysfunction due to atrial fibrillation in patients initially believed to have idiopathic dilated cardiomyopathy. *Am J Cardiol.* 1992;69:1570-3.
11. Wang TJ, Larson MG, Levy D, et al. Temporal relations of atrial fibrillation and congestive heart failure and their joint influence on mortality: the Framingham Heart Study. *Circulation.* 2003; 107:2920-5.
12. Humphries KH, Jackevicius C, Gong Y, et al. Population rates of hospitalization for atrial fibrillation/flutter in Canada. *Can J Cardiol.* 2004;20:869-76.
13. Stewart S, Hart CL, Hole DJ, McMurray JJ. Population prevalence, incidence, and predictors of atrial fibrillation in the Renfrew/Paisley study. *Heart.* 2001;86:516-21.
14. Geraci SA, Haan CK. Effect of beta blockers after coronary artery bypass in postinfarct patients: what can we learn from available literature? *Ann Thorac Surg.* 2002;74:1727-32.
15. Effect of enalapril on survival in patients with reduced left ventricular ejection fractions and congestive heart failure. The SOLVD Investigators. *N Engl J Med.* 1991;325:293-302.
16. Ferguson TB Jr, Coombs LP, Peterson ED. Preoperative beta-blocker use and mortality and morbidity following CABG surgery in North America. *JAMA.* 2002;287:2221-7.
17. Gajewski J, Singer RB. Mortality in an insured population with atrial fibrillation. *JAMA.* 1981;245:1540-4.
18. Franz MR, Karasik PL, Li C, Moubarak J, Chavez M. Electrical remodeling of the human atrium: similar effects in patients with chronic atrial fibrillation and atrial flutter. *J Am Coll Cardiol.* 1997; 30:1785-92.
19. Allessie MA. Atrial fibrillation-induced electrical remodeling in humans: what is the next step? *Cardiovasc Res.* 1999;44:10-2.
20. Chua YL, Schaff HV, Orszulak TA, Morris JJ. Outcome of mitral valve repair in patients with preoperative atrial fibrillation. Should the maze procedure be combined with mitral valvuloplasty? *J Thorac Cardiovasc Surg.* 1994;107:408-15.
21. Savelieva I, Camm AJ. Clinical relevance of silent atrial fibrillation: prevalence, prognosis, quality of life, and management. *J Interv Card Electrophysiol.* 2000;4:369-82.

### Notice of Correction

Shigemura N, Akashi A, Funaki S, Nakagiri T, Inoue M, Sawabata N, et al. Long-term outcomes after a variety of video-assisted thoracoscopic lobectomy approaches for clinical stage IA lung cancer: A multi-institutional study. *J Thorac Cardiovasc Surg.* 2006;132:507-12.

The authors report the following error in data analysis: [Figure 3](#) shows the postoperative physical activity of the patients using an accelerometer to quantify their recovery from surgery. The data pertaining to all patients in one of the participant hospitals who received the a-VATS approach are inaccurate. All a-VATS results have been removed from the revised [Figure 3](#). As a result, the authors can no longer conclude that a significant difference exists in the recovery times between the a-VATS and c-VATS groups. The significant difference in the recovery times between the c-VATS and open thoracotomy groups remains valid. All other results from the study remain accurate.



**Revised Figure 3.** Comparison of the postoperative physical ability as determined by Active Tracer and expressed as the percentage of the preoperative 24-hour value. Time points included are before surgery and 0, 1, 2, 3, 4, 5, and 6 days postoperatively. Each value represents the mean  $\pm$  standard error at each time point. \* $P < .05$  for complete VATS (c-VATS) versus open thoracotomy on postoperative days 2 to 5.